



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}\int \cos \theta^2 d\theta &= \theta \sum_{t=0}^{\infty} 2^t \prod_{s=1}^{s=t} \frac{1}{2s+1} \theta^{2t} \cos \left(t \frac{\pi}{2} - \theta^2 \right) \\ &= \cos \theta^2 \left(\theta - \frac{2^2}{1 \cdot 3 \cdot 5} \theta^5 + \frac{2^4}{1 \cdot 3 \cdot 5 \cdot 7 \cdot 9} \theta^9 - + \dots \right) \\ &\quad + \sin \theta^2 \left(\frac{2}{1 \cdot 3} \theta^3 - \frac{2^3}{1 \cdot 3 \cdot 5 \cdot 7} \theta^7 + - \dots \right).\end{aligned}$$

Two interesting special cases deserve mention.

When the exponent of the coefficient is less by unity than that of the argument of the function, the integral assumes a known form; and the value of the other integral becomes known. Hence, expressions of the type $\theta^{n-1} \sin \theta^n d\theta$ are directly integrable in finite form by these formulæ. If in addition n takes the special form $1/r$, the integral of $\sin \theta^{1/r} d\theta$ is found, that is, the integral of sine or cosine of any root of a variable is found in finite form.

Thus, taking the first of the four formulæ, making $m = n - 1$, and transforming a little; and after this, making $n = 1/r$, we obtain the formulæ:

$$\begin{aligned}\int \theta^{rn-1} \cos \left(r \frac{\pi}{2} - \theta^n \right) d\theta &= -n^{-1} \prod_{s=r-1}^{s=1} s \cdot \sum_{t=1}^{t=r} \prod_{s=1}^{s=t-1} s^{-1} \cdot \theta^{(t-1)n} \sin \left(t \frac{\pi}{2} - \theta^n \right), \\ \int \cos \left(r \frac{\pi}{2} - \theta^{1/r} \right) d\theta &= -r \prod_{s=1}^{s=r-1} s \cdot \sum_{t=1}^{t=r} \prod_{s=1}^{s=t-1} s^{-1} \cdot \theta^{(t-1)/r} \sin \left(t \frac{\pi}{2} - \theta^{1/r} \right).\end{aligned}$$

As examples, in these formulas making $n = 2$, $r = 4$, we have

$$\begin{aligned}\therefore \int \theta^7 \cos \theta^2 d\theta &= -3 \left[\cos \theta^2 \left(1 - \frac{\theta^4}{2} \right) + \sin \theta^2 \left(\theta^2 - \frac{\theta^6}{3!} \right) \right], \\ \int \cos \theta^{1/4} &= -4! \left[\cos \theta^{1/4} \left(1 - \frac{\theta^{1/2}}{2} \right) + \sin \theta^{1/4} \left(\theta^{1/4} - \frac{\theta^{3/4}}{3!} \right) \right].\end{aligned}$$

DISCUSSIONS.

I. RELATING TO A CURVE WITH UNUSUAL PROPERTIES.

By JOS. B. REYNOLDS, Lehigh University.

The curve

$$y = \frac{a^2 x}{x^2 + a^2}$$

presents some unusual properties. Considering the part of the curve to the right of the y -axis (see the figure) we have for the area between it and the x -axis

$$A = \int_0^{\infty} y dx = a^2 \int_0^{\infty} \frac{x dx}{x^2 + a^2} = \frac{a^2}{2} \log (x^2 + a^2) \Big|_0^{\infty} = \infty.$$


Yet the volume generated by revolving this area about the x -axis is

$$V = \pi \int_0^{\infty} y^2 dx = \pi a^4 \int_0^{\infty} \frac{x^2 dx}{(x^2 + a^2)^2} = \frac{\pi^2 a^3}{4};$$

that is, the revolution of an infinite area gives a finite volume. Again, the ordinate of the center of gravity of this area is

$$\bar{y} = \frac{\int_0^\infty dx \int_0^{a^2x/(x^2+a^2)} y dy}{\int_0^\infty dx \int_0^{a^2x/(x^2+a^2)} dy} = \frac{\frac{\pi a^3}{8}}{\infty} = 0;$$

that is, although the area lies entirely above the x -axis and is indefinitely large,



its center of gravity is indefinitely close to that axis.

By the theorem of Pappus we have

$$2\pi\bar{y}A = V,$$

whence we have an indeterminate form evaluated


$$2\pi(0)\infty = \frac{\pi^2 a^3}{4}.$$

II. RELATING TO A DEMONSTRATION OF A GEOMETRICAL THEOREM.

By WILLIAM E. HEAL, Washington, D. C.

If the bisectors of two angles of a triangle are equal, the triangle is isosceles.

This is a very celebrated theorem and has been demonstrated in many ways. It was proposed in the MONTHLY as problem 42 and several demonstrations were published in Vol. II, pp. 157 and 189-191. Of these proofs all but one were by indirect methods; that is, by use of the *reductio ad absurdum*. The following direct method of proof was communicated to the writer some months ago by Dr. Artemas Martin of the U. S. Coast Survey and seems to be eminently worthy of preservation.



If $BD = CE$ we are to prove that $AB = AC$.

We have by a well-known theorem

$$AB \times BC = (AE + BE)BC = BD^2 + AD \times DC, \quad (1)$$

$$AC \times BC = (AD + DC)BC = CE^2 + AE \times EB. \quad (2)$$

If $BD = CE$ these become

$$(AE + BE)BC = BD^2 + AD \times DC, \quad (3)$$

$$(AD + DC)BC = BD^2 + AE \times EB. \quad (4)$$